

COMMISSION 27 OF THE I. A. U.
 INFORMATION BULLETIN ON VARIABLE STARS

Number 1368

Konkoly Observatory
 Budapest
 1977 December 7

PHOTOELECTRIC MINIMA OF U Oph, AB And AND X Tri

Following are the results of observations made at the Ankara University Observatory with a 30 cm Maksutov telescope, an EMI 6256 S photomultiplier tube and standard B, V filters:

Star	Hel.Min.J.D.	m.e.	Min.	O-C
U Oph	2443319.3964	±.0004	I	+ .0026
	34.4917	.0012	I	.0018
	35.3291	.0005	II	.0005
	40.3622	.0003	II	.0016
	45.3950	.0008	II	.0024
	56.2978	.0005	I	.0024
	61.3304	.0006	I	.0030
	66.3629	.0005	I	.0034
AB And	2443369.3904	±.0015	I	+ .0022
	70.3873	.0004	I	.0035
	71.3827	.0005	I	.0032
	73.3744	.0003	I	.0035
	75.3653	.0002	I	.0031
X Tri	2443379.5316	±.0020	II	+ .0096
	80.5006	.0012	II	.0070
	98.4753	.0006	I	.0085

SAC 1977 formula $\text{Min.I.} = \text{J.D. } 2436727.424 + 1^d 6773460 \cdot \text{E}$.
 Quester's formula $\text{Min.I.} = \text{J.D. } 2436109.57928 + 0.33189215 \cdot \text{E}$ (IBVS 190)
 and SAC 1977 formula $\text{Min.I.} = \text{J.D. } 2440984.2205 + 0.9715277 \cdot \text{E}$
 were used, respectively, in the calculations of O-C's.

ZEKİ TÜFEKÇİOĞLU
 Ankara University Observatory
 A.Ü. Fen Fakültesi
 Ankara-Turkey