

COMMISSION 27 OF THE I. A. U.
INFORMATION BULLETIN ON VARIABLE STARS

Number 2159

Konkoly Observatory
Budapest
1982 June 14
HU ISSN 0374-0676

TIMES OF MINIMA FOR EIGHT ECLIPSING BINARIES

Thirteen times of minima for eight eclipsing variables have been determined photoelectrically with the 30 cm Maksutov telescope of the University of Ankara, using an uncooled EMI 6256S photomultiplier tube and a dc chart recorder.

| <u>Binary</u> | <u>HJD</u> | <u>m. e.</u> |
|---------------|-----------------------|--------------|
| ST Aqr | 2444845.3898 ± 0.0004 | |
| RZ Com | 694.3712 ± 0.0006 | |
| | 695.3868 ± 0.0003 | |
| RW CrB | 780.4044 ± 0.0002 | |
| | 783.3109 ± 0.0010 | |
| V836 Cyg | 840.4236 ± 0.0003 | |
| | 842.3839 ± 0.0002 | |
| V1073 Cyg | 783.4139 ± 0.0005 | |
| | 790.4899 ± 0.0005 | |
| TZ Lyr | 784.4082 ± 0.0005 | |
| BB Peg | 812.5022 ± 0.0003 | |
| DI Peg | 843.4272 ± 0.0002 | |
| | 848.4102 ± 0.0002 | |

Each minimum time given in the table is the mean of two determinations through b and v filters. The timings were made by the method of Kwee and Van Woerden (1956).

Supports by TUBITAK (Scientific and Technical Research Council

of Turkey) and US National Science Foundation through a grant to Prof. T.J. Herczeg of the University of Oklahoma are gratefully acknowledged.

E. DERMAN, N.YILMAZ, S.ENGİN
Z. ASLAN, C.AYDIN, Z.TÜFEKÇİOĞLU

University Observatory
A.Ü. Fen Fakültesi
Ankara, Turkey

Reference:

K.K.Kwee and H. Van Woerden, Bull. Astron. Neth. 12, 327, 1956