


### HD 209775: A NEW $\delta$ SCUTI VARIABLE

HENRY, G.W.; HENRY, S.M.

Center of Excellence in Information Systems, Tennessee State University, 330 10th Ave. North, Nashville, Tennessee 37203, U.S.A., e-mail: henry@schwab.tsuniv.edu, henrysm@galileo.tsuniv.edu


In the course of making photometric observations of planetary transits in HD 209458 (Henry et al. 1999; 2000), we found one of our comparison stars, HD 209775, to be a new small-amplitude variable. Very little has been published about HD 209775; Olsen (1980) estimated a spectral type of Am/p based on his Strömngren *uvby* photometry and the assumption that his Strömngren indices were not contaminated by duplicity. Fehrenbach & Burnage (1982) found the radial velocity to be constant to  $2.6 \text{ km s}^{-1}$  in six measurements. The *HIPPARCOS* catalogue classifies the star as photometrically constant and gives  $V = 7.56$ ,  $B - V = 0.33$ , and  $\pi = 10.17 \text{ mas}$  (Perryman et al. 1997). Combined with the effective temperature calibration of Flower (1996), the *HIPPARCOS* results imply an absolute magnitude of 2.50 and a spectral type of F0 V.



**Figure 1.** Johnson *B* photometry over two nights acquired with a 0.40 m APT showing HD 209775 to be a new low-amplitude, short-period variable star.

Figure 1 plots our Johnson *B* photometry acquired with the T3 0.40 m automatic photoelectric telescope (APT) located at Fairborn Observatory in southern Arizona. We

used HD 209458 ( $V = 7.65$ ,  $B - V = 0.594$ , G0 V) as a comparison star to compute the differential magnitudes since it has been shown to be constant to 0.0017 mag or better except during transits (Henry et al. 2000). The observations were corrected for differential extinction and transformed to the standard Johnson system. Details on the operation of the APT and the reduction and precision of the data can be found in Henry (1995).



**Figure 2.** Periodogram analysis of the  $B$  data gives a period of  $0.0608 \pm 0.0002$  d and a full amplitude of  $0.012 \pm 0.001$  mag.

The results of periodogram analysis of the  $B$  observations from Figure 1 are shown in Figure 2. We find a period of  $0.0608 \pm 0.0002$  d. A least-squares sine fit to the  $B$  data at this period gives a full amplitude of  $0.012 \pm 0.001$  mag. From the  $V$  observations, we find a period of  $0.0607 \pm 0.0002$  d and an amplitude of  $0.007 \pm 0.001$  mag. The star lies in the region of the HR diagram where the  $\delta$  Scuti and the  $\gamma$  Doradus variables overlap (Handler 1999). Its short period places it in the  $\delta$  Scuti class.

Astronomy with automated telescopes at Tennessee State University is supported through NASA grants NCC5-96 and NCC5-228, which funds the Center for Automated Space Science, and NSF grant HRD-9706268, which funds the Center for System Science Research.

#### References:

- Fehrenbach, C., & Burnage, R., 1982, A&AS, 49, 483  
 Flower, P. J., 1996, ApJ, 469, 355  
 Handler, G., 1999, MNRAS, 309, L19  
 Henry, G. W., 1995, in Robotic Telescopes: Current Capabilities, Present Developments, and Future Prospects for Automated Astronomy ASP Conf. Ser. No. 79, eds. G. W. Henry and J. A. Eaton (ASP: San Francisco), p. 44  
 Henry, G. W., Marcy, G. W., Butler, R. P., & Vogt, S. S., 1999, IAU Circular No. 7307  
 Henry, G. W., Marcy, G. W., Butler, R. P., & Vogt, S. S., 2000, ApJ, 529, L41  
 Olsen, E. H., 1980, A&AS, 39, 205  
 Perryman, M. A. C., et al., 1997, The Hipparcos and Tycho Catalogues (ESA: The Netherlands)