# COMMISSIONS 27 AND 42 OF THE IAU INFORMATION BULLETIN ON VARIABLE STARS

Number 5065

Konkoly Observatory Budapest 25 April 2001

 $HU\ ISSN\ 0374-0676$ 

### NSV 03007 IS AN EW ECLIPSING BINARY SYSTEM

GARCIA-MELENDO, E.<sup>1</sup>; SANCHEZ-BAJO, F.<sup>2</sup>

 $^1$ Esteve Duran Observatory Foundation, Montsen<br/>y46– Urb. El Montanya, 08553 Seva, Barcelona, Spain, e-mail: duran<br/>obs@astrogea.org

 $^2$ Departamento de Electrónica e Ingenieria Electromecánica, Escuela de Ingenierias Industriales, Universidad de Extremadura, Ctra de Elvas s/n, 06071 Badajoz, Spain e-mail: fsanbajo@unex.es

Name of the object:		
$NSV \ 03007 = GSC \ 3376 \ 287 = HV \ 07649 = CSV \ 000766$		
Equatorial coordinates:		Equinox:
<b>R.A.</b> = $20^{h}22^{m}10^{s}$ <b>DEC.</b> = $+31^{\circ}15'12''$		2000.0
Observatory and telescope:		
Esteve Duran Observatory, 0.6-m Cassegrain telescope		
Detector:	CCD	
$\mathbf{T}$ : $\mathbf{I}_{\mathbf{r}}$	Ι.Z	
Filter(s):	V	
Comparison star(s):	GSC 3376_1091	
Check star(s):	GSC 3376_254	
Transformed to a standard system: No		
· · · · · · · · · · · · · · · · · · ·		
Availability of the data:		
Upon request		
Type of variability:	EW	




Figure 1. Folded light-curve of NSV 03007

#### **Remarks:**

The variability of NSV 03007 was announced by Boyd in 1937, and listed in the NSV catalogue (Kholopov, 1982) as an RR Lyr variable with a photographic magnitude variation between 13.7 and 14.3. The star, observed during a NSV survey searching for faint variables, was successfully monitored for 6 nights from October 2, 2000 to March 18, 2001. Observations show that this object is actually an EW eclipsing binary system with a period of about  $0^{d}385$  (Figure 1). The star fades  $0^{m}44$  during minimum I and 0.40 during the secondary eclipse. Taking as a reference the Vmagnitude of the check star derived from Tycho observations (ESA, 1997), our photometry indicates that the minimum V magnitude for NSV 03007 is of 13.17. Although some fundamental observational data about this star are still unknown, a preliminary analysis using the Wilson–Devinney code (Wilson, 1998) suggests that the mass ratio between the two components could be as small as 0.2. The following ephemeris was computed:

> Min. I = HJD 2451966.464 +  $0.384758 \times E$ ,  $\pm\,0.002\,\pm 0.000020$

in addition to the two secondary minimum timings: HJD 2451960.498 and 2451962.422.

#### References:

Boyd, C. D., 1937, Harvard Bulletin, No. 905 ESA, 1997, The Hipparcos and Tycho Catalogues, ESA SP-1200 Kholopov, P. N., editor, 1982, New Catalogue of Suspected Variable Stars, Moscow Wilson, R. E., 1998, Computing Binary Star Observables (Reference Manual to the

## ERRATUM FOR IBVS 5065

The coordinates for NSV 03007 given in IBVS 5065 are in error. The actual ones (2000), according to its identification with GSC 3376-0287 and SIMBAD database are:

 $\begin{aligned} {\rm R.A.} &= ~06^{\rm h}32^{\rm m}46^{\rm s}\!.2 \\ {\rm Dec.} &= ~+46^\circ23'32'\!\!.82\,. \end{aligned}$ 

Enríque García-Melendo