COMMISSIONS 27 AND 42 OF THE IAU INFORMATION BULLETIN ON VARIABLE STARS

Number 5170

Konkoly Observatory Budapest 13 September 2001 *HU ISSN 0374 - 0676*

UZ CVn: A CENTURY OF PERIOD INCREASE

VANDENBROERE, JACQUELINE¹; BERTHOLD, THOMAS^{2,3}

¹ Group Européen d'Observations Stellaires (GEOS), 23 Parc de Levesville, F–28300 Bailleau l'Evëque, France, e-mail: j.vandenbroere@skynet.be

 2 Bundesdeutsche Arbeitsgemeinschaft für Veränderliche Sterne e.V. (BAV), Munsterdamm 90,

³ Bruno-H.-Bürgel-Sternwarte Hartha, Töpelstr. 43, D–04746 Hartha, Germany, e-mail: tb.stwhartha@web.de

UZ CVn (= BV 96 = HIP 61029 = GSC 3018 255, $12^{h}30^{m}27^{s}7$, $+40^{\circ}30'31''_{.9}$, 2000.0), was discovered by Kippenhahn (1955) and found to be a pulsating star of RRab type by Strohmeier and Knigge (1961). Their elements, listed in the GCVS (Kholopov, 1985),

$$Max. = HJD \ 2426427.3806 + 0^{d}6977829 \times E$$
(1)

were derived from photographic plates They are obviously no longer valuable because the period has changed continuously. In order to study the evolution of the period of UZ CVn during the last century, we have gathered all the instants of maximum light published in the literature and we have also performed our own measurements.

There are only 13 photoelectric measurements in V and R made between March 1990 and December 1991 resulting in the determination of one instant of maximum (Schmidt et al., 1995). The 127 CCD transits accepted from the Hipparcos satellite measurements (1990–1993) were studied by Fernley et al. (1998) who obtained a period of 0^d.697783 with a very scattered light curve. We took into account the epoch of the Hipparcos catalogue only. The photographic data are very scattered instants of bright light obtained from the inspection of sky patrol plates. The first 59 times of maximum found from Bamberg and Sonneberg plates taken between 1931 and 1960, including 5 instants published before by Filatov (1960) and 38 instants derived by Döppner from Sonneberg plates, were published by Strohmeier and Knigge (1961), establishing ephemeris (1). A re-inspection of the Sonneberg plates done by one of us (T.B.) has evidenced the timings assigned to Döppner to be in fact geocentric! They have been corrected for further analysis. Later on, a further set of 80 instants of bright light was published by Strohmeier and Bauernfeind (1968) as a result of the investigation of Harvard photographic plates taken between 1901 and 1953.

A GEOS team made 26 photoelectric measurements of UZ CVn in the B and V filters of the Geneva system at the Jungfraujoch observatory during two nights in January 1997. A new time of maximum could be determined. Seven additional measurements were obtained at the same observatory in 1998 (see Figure 1). Two visual observers, J.-P. Verrot and J. Vandenbroere, determined 20 further instants of maximum from their estimates made between 1994 and 2001. To close the remaining gap in the data between 1960 and 1990, T.B. has used 554 Sonneberg Observatory sky patrol plates taken between

D–12169 Berlin, Germany



Figure 1. V lightcurve of UZ CVn according to ephemeris (2)



Figure 2. O - C diagram of UZ CVn according ephemeris (2). The symbols refer to the kind of observation: \times (visual), + (photographic), * (photographic normal maxima), \circ (HIPPARCOS), • (photoelectric)

1959 and 1993. From 69 newly found instants, 7 normal maxima in consecutive intervals were derived for further analysis.

Taking into account all the available material, we are able to document the behaviour of the period of an RR Lyrae star over a whole century. The complete list of all the observed times of maximum is available from the IBVS website as file 5170-t1.txt. A linear least-squares fit, made with 163 instants of maximum, consisting of two photoelectric instants (w = 10), seven photographic normal maxima (w = 4), one Hipparcos and 20 visual instants (w = 3) and 133 photographic instants (w = 1), covering the years from 1901 to 2001, has yielded the following ephemeris:

$$Max. = HJD \ 2415423.9927 + 0^{d}.69778714 \times E. \\ \pm 74 \qquad \pm 47$$
(2)

As Figure 2 points out, the trend of the O - C values can be represented either by an abrupt period change around epoch 28000 or by a parabolic fit.

From JD 2415400 (approx.) to JD 2435000 (approx.):

$$Max. = HJD \ 2415424.1137 + 0\overset{d}{.}69777993 \times E. \\ \pm 120 \qquad \pm 66$$
(3)

From JD 2435000 (approx.) to JD 2452100 (approx.):

$$Max. = HJD \ 2450460.6095 + 0^{d}.69779191 \times E. \\ \pm 67 \qquad \pm 15$$
(4)

Alternatively, the quadratic least squares fit yields the following elements:

$$Max. = HJD \ 2415424.1453 + 0^{d}.69777362 \times E + 2.19 \times 10^{-10} \times E^{2}. \\ \pm 91 \qquad \pm 69 \qquad \pm 11$$
(5)

Assuming the last case, the period of UZ CVn has been established to have increased by a constant rate of $dP = 6.28 \times 10^{-10}$ per day during the last century and thus has increased by 1.98s in the same time. Such rates are found to be typical in numerous cases among RR Lyrae variables.

We want to acknowledge M. Dumont (GEOS) and L. Zimmerman (Cercle Astronomique de Bruxelles) for discussion and Peter Kroll (Sonneberg Observatory) for the use of the plate archive. This research made use of the SIMBAD database operated by the CDS at Strasbourg.

References:

Fernley, J., et al., 1998, Astron. Astrophys., 330, 515
Filatov, G.S., 1960, Astron. Circ., 215, 17
Kholopov, P.N., et al., 1985, GCVS, IVth edition, Nauka, Moscow Kippenhahn, R., 1955, Kl. Ver. Bamberg, 11
Schmidt, E.G., et al., 1995, Astron. Journ., 109, 1239
Strohmeier, W., and Bauernfeind, H., 1968, Bamberg Ver., VII, 77
Strohmeier, W., and Knigge, R., 1961, Bamberg Ver., V, 11