

COMMISSIONS 27 AND 42 OF THE IAU  
 INFORMATION BULLETIN ON VARIABLE STARS

Number 5211

Konkoly Observatory  
 Budapest  
 13 December 2001

*HU ISSN 0374 – 0676*

**PHOTOMETRIC VARIABILITY OF THE STARS  
 USNO 0900-17903132, HD191674 AND HD 191616**


ROBB<sup>1</sup>, R. M.; LINDNER, J. V.; STERN, L. A.; LINDNER, T. H.; BALAM<sup>1,2</sup>, D. D.; CLEM, J. L.

<sup>1</sup> Guest User, Canadian Astronomy Data Centre, which is operated by the Herzberg Institute of Astrophysics, National Research Council of Canada

<sup>2</sup> Guest Observer, Dominion Astrophysical Observatory, which is operated by the Herzberg Institute of Astrophysics, National Research Council of Canada

Climenhaga Observatory, Dept. of Physics and Astronomy, University of Victoria, Victoria, BC, Canada, V8W 3P6, Internet: robb@uvic.ca

As part of a continuing search for photometric variations in stars which are known X-ray sources, we observed HD 191616 (=GSC 0503-0700). This star, also known as RXJ201043+045449, was discovered to be a source of X-rays by the ROSAT satellite (Voges et al. 1999). The Tycho catalog (ESA 1997) reported  $V_T=9^m497$  and  $B_T=10^m724$  for HD 191616 and  $V_T=9^m000$  and  $B_T=10^m238$  for one of the field stars, HD 191674 (=GSC 0503-1409), both consistent with a spectral type of approximately K0. The reported parallaxes favor dwarf luminosity class but do not exclude giant status.



**Figure 1.** Finder chart labeled with the GSC identification numbers from region 0503.

The field of stars observed with the automated 0.5m telescope of the Climenhaga Observatory at the University of Victoria is shown in Figure 1. The data were reduced in a fashion similar to that described in Robb and Greimel (1999). Table 1 lists the stars' identification numbers, coordinates (J2000) and magnitudes from the Hubble Space


Table 1: Stars observed in the field of HD 191616=GSC 0503-0700 and HD 191674=GSC 0503-1409

Star Id	R.A. J2000	Dec. J2000	GSC Mag.	$\Delta R_c$ Mag.	Std Dev Between	Std Dev Within	( $R - I$ )
HD 191616	20 <sup>h</sup> 10 <sup>m</sup> 44 <sup>s</sup>	4°54'50"	8.8	0.490	.022	.004	0.4
GSC 0503-1521	20 <sup>h</sup> 10 <sup>m</sup> 51 <sup>s</sup>	4°56'57"	8.6	-	-	-	0.4
HD 191674	20 <sup>h</sup> 11 <sup>m</sup> 03 <sup>s</sup>	4°55'10"	9.3	-.222	.025	.003	0.4
GSC 0503-1318	20 <sup>h</sup> 10 <sup>m</sup> 38 <sup>s</sup>	4°54'19"	10.0	1.447	.003	.003	0.4
GSC 0503-388	20 <sup>h</sup> 10 <sup>m</sup> 35 <sup>s</sup>	4°54'55"	11.9	3.428	.004	.007	0.2
GSC 0503-328	20 <sup>h</sup> 10 <sup>m</sup> 43 <sup>s</sup>	4°56'09"	11.7	3.060	.004	.009	0.4
USNO 900-17903132	20 <sup>h</sup> 10 <sup>m</sup> 43 <sup>s</sup>	4°55'51"	13.9	4.787	.157	.036	2.3
GSC 0503-802	20 <sup>h</sup> 10 <sup>m</sup> 54 <sup>s</sup>	4°56'45"	12.4	3.474	.002	.007	0.3
GSC 0503-2279	20 <sup>h</sup> 10 <sup>m</sup> 47 <sup>s</sup>	4°58'44"	13.1	4.579	.011	.019	0.2


Telescope Guide Star Catalog (GSC) (Jenkner et al., 1990). Observations were made using filters closely matching the Cousins R and I bands (Cousins 1981). The thirty-two Julian Dates of observation ( $-2450000$ ) are 2121, 2122, 2129-2138, 2150, 2151, 2157-2166, 2168, 2169, 2175, 2176, 2179, 2180, 2182 and 2184.

Our differential  $\Delta R_c$  magnitudes are calculated in the sense of the star minus GSC 0503-1521. Brightness variations during a night were measured by the standard deviation of the differential magnitudes and are listed for the most photometric night in the last column as “Std Dev Within”. For each star the mean of the nightly means is shown as  $\Delta R_c$  in Table 1. The standard deviation of the nightly means is a measure of the night to night variations and is called “Std Dev Between” in Table 1.

The “Std Dev Between” for stars [GSC 0503-1318 - GSC 0503-1521] is 0.003 magnitudes. This excellent photometry shows that night to night variations in either of these stars must be less than a few millimagnitudes. We observed no significant variations in these stars in plots of the individual nights’ data and a “Std Dev Within” one night of 0.003 sets an upper limit on variations of an hourly timescale.

Figure 2.  $\Delta R_c$  filtered light curve of USNO 0900-17903132 for summer 2001

In Figure 2 we have plotted the nightly mean and standard deviation of the  $\Delta R_c$  brightness of the star USNO 0900-17903132. Obviously it is a variable star with a timescale of at least a hundred days and from our  $(R - I)_c$  we estimate a spectral type of late M. We expect this star to be a long period variable and urge observers to monitor its brightness and discover its amplitude and periodicity, if any.




**Figure 3.** Periodogram for HD 191674 in 2001

The star HD 191674 had obvious night to night variations. The root-mean-square (RMS) deviation of a data point from a sine curve as a function of period is shown in Figure 3. From this periodogram we find the ephemeris to be:

$$\text{HJD of Maximum Brightness} = 2452118^{\text{d}}.3(9) + 5^{\text{d}}.21(15) \times E.$$

where the uncertainties in the final digit are given in brackets and the RMS error of the fit is less than 0.01 magnitudes. The 3685 differential  $\Delta R_c$  magnitudes phased at this period are plotted in Figure 4 with different symbols for each cycle.




**Figure 4.**  $R_c$  filtered light curve of HD 191674

There are significant differences from cycle to cycle but doubling the period did not improve the fit. The  $(R - I)_c$  is consistent with the Tycho  $B_T$  and  $V_T$  measurements indicating that the star is approximately K0 spectral type. Therefore we postulate that HD 191674 is a late type spotted dwarf rotating with a  $5^{\text{d}}.21$  period.


Our observations of HD 191616 are plotted in Figure 5. with the error bars set to the standard deviation of the points during each night. The error of the nightly mean then would be about ten times smaller. Obviously HD 191616 is also a variable star, so Period98 (Sperl 1998) was used to find the period, amplitude and shape of the variations. The best model light curve found is of the form:

$$\Delta R = 0.4611 + 0.0554 * \sin(2\pi(\text{JD} - 2112.0)/18.4638) + 0.0198 * \sin(2\pi(\text{JD} - 2113.572)/9.2319)$$



**Figure 5.**  $\Delta R_c$  filtered light curve of HD 191616 for 2001 with model curve overplotted

Note that the second period was set to exactly half the dominant period, to modify the shape of the curve to best match the asymmetry in the distribution of the spots on the surface of the star. Spectroscopic observations made with the DAO 1.8m telescope were used to determine the spectral class to be K3, consistent with the  $(R - I)_c$  and Tycho measurements. The spectrum displayed in figure 5 clearly shows the  $H\alpha$  emission line.



**Figure 6.** Spectrum of HD 191616 showing the  $H\alpha$  emission line at  $6563\text{\AA}$ .

HD 191616 is an X-ray source, a K3 spectral type star with  $H\alpha$  emission, photometric variations with an  $18^d.5 \pm .4$  period and cycle to cycle differences typical of BY Dra stars. Photometric observations should be continued to monitor for flares, changes in the spot distribution and to determine the period more precisely.

#### References:

- Cousins, A.W., 1981, SAAO Circ., **6**, 4  
 ESA, 1997, The Hipparcos and Tycho Catalogues, ESA SP-1200  
 Jenkner, H., Lasker, B., Sturch, C., McLean, B, Shara, M., Russell, J., 1990, *AJ*, **99**, 2082  
 Robb, R.M. and Greimel, R., 1999, *ASP Conf. Ser.*, **189**, 198  
 Sperl, M., 1998, Master's Thesis, Institut für Astronomie, Universität Wien  
 Voges, W., et al., 1999, *A&A*, **349**, 389