COMMISSIONS G1 AND G4 OF THE IAU INFORMATION BULLETIN ON VARIABLE STARS

Volume 63 Number 6259

DOI: 10.22444/IBVS.6259

Konkoly Observatory Budapest 29 January 2019 *HU ISSN 0374 - 0676*

DETECTION OF A δ SCUTI–TYPE PULSATING COMPONENT IN THE DETACHED ECLIPSING BINARY SYSTEM TU CMa

MKRTICHIAN, D.E.

National Astronomical Research Institute of Thailand, 260 Moo 4, T. Donkaew, A. Maerim, Chiangmai, 50180 Thailand. e-mail: davidmkrt@gmail.com

Abstract

We report the detection of 30.5 min low-amplitude (A=0.003 mag) δ Scuti-type pulsations in an A5V-A7V type component of the detached eclipsing binary system TU CMa.

TU CMa is a 1.127803854-day (Haans et al. 2015) detached eclipsing binary system with A5V-A7V primary and F8V-G0V secondary components (Garces et al. 2017). It was included in our program to search for pulsating components that have the primary component lying inside the instability strip and hence can be potentially pulsating.

Visual inspection of the SWASP data of TU CMa taken from their archive¹ revealed about 0.02-day short-period, low-amplitude light variations. For the safe detection of possible pulsations, we selected the best quality nights for TU CMa from the SWASP data, namely HJD 2454105, 2454131, 2454132, 2454133, 2454134, 2454135, 2454433, 2454434, 2454436, 2454456, 2454462 and 2454485. The pulsation variations were searched for in the out-of-eclipse parts of the light curves after removal of slow orbital variations using low order polynomial fits. For the period search, we used the PERIOD04 software (Lenz & Breger, 2005) based on a Discrete Fourier Transform (DFT) analysis.

The DFT amplitude-frequency spectrum of the TU CMa residual data is shown in Figure 1. We detected a clear signal at 47.3197 ± 0.0002 c/d (P = 30.5 min) with an amplitude of 0.0038 mag. The phased light curve binned into 20-phase intervals is shown in Figure 2.

Using the mass M=1.761±0.012 M_{\odot} and the radius R= 1.553±0.002 R_{\odot} for the primary component from Garces et al. (2017) we calculated the mean density of the pulsating component as $\rho/\rho_{\odot}=0.4702$. The calculated pulsation constant for the discovered 30.5 min (P = 0.021 day) pulsation mode, $Q = P_{\sqrt{\rho_*/\rho_{\odot}}} = 0.014$, corresponds to a fourth or fifth overtone low degree ($\ell = 0 - 3$) mode.

Conclusion: We report the detection of a 30.5 min low amplitude (A=0.003 mag) δ Scuti-type pulsation in an A5V-A7V type component of the detached eclipsing binary system TU CMa. The calculated pulsation constant corresponds to pulsations in the 4-5th overtone low-degree mode (Fitch, 1981). The parameters of the binary system and

¹https://wasp.cerit-sc.cz



Figure 1. The DFT amplitude spectrum of the primary A5V-A7V component. The dominant peak is at 47.3197 ± 0.0002 c/d.



Figure 2. The phase-binned pulsation light variations of TU CMa. The phase of the maximum light corresponds to HJD 2454107.9776.

components of TU CMa are accurately determined and are good input parameters for theoretical pulsational modelling. This binary system can be a good target for further more accurate and detailed photometric observations of pulsations in order to detect a low-amplitude pulsation spectrum, for the eclipse mode identification of the dominant mode and for comparison with theoretical pulsation models.

Acknowledgements: I acknowledge this work as part of the research activity supported by the National Astronomical Research Institute of Thailand (NARIT), Ministry of Science and Technology of Thailand.

This paper makes use of data from the DR1 of the WASP data (Butters et al. 2010) as provided by the WASP consortium, and the computing and storage facilities at the CERIT Scientific Cloud, reg. no. CZ.1.05/3.2.00/08.0144 which is operated by Masaryk University, Czech Republic (Paunzen et al., 2014).

References:

Butters, O. W., et al., 2010, A&A,520, L10 DOI
Fitch, W. S., 1981, ApJ, 249, 218 DOI
Garcés, J. L., Mennickent, R. E., & Zharikov, S, 2017, PASP 129, 044203 DOI
Haans, G. K., et al., 2015, PKAS, 30, 205 DOI
Lenz, P., Breger M., 2005, Communications in Asteroseismology, 146, 53 DOI
Paunzen, E., Kuba, M., West, R.G., Zejda, M., 2014, IBVS, 6090